

On Rigidity of Roe Algebras

Ján Špakula and Rufus Willett

October 10, 2011

Abstract

Roe algebras are C^* -algebras built using large-scale (or ‘coarse’) aspects of a metric space (X, d) . In the special case that $X = \Gamma$ is a finitely generated group and d is a word metric, the simplest Roe algebra associated to (Γ, d) is isomorphic to the crossed product C^* -algebra $l^\infty(\Gamma) \rtimes_r \Gamma$.

Roe algebras are ‘coarse invariants’, in the sense that if X and Y are coarsely equivalent metric spaces, then their Roe algebras are isomorphic. Motivated in part by the coarse Baum-Connes conjecture, we ask if there is a converse to the above statement: that is, if X and Y are metric spaces with isomorphic Roe algebras, must X and Y be coarsely equivalent? We show that for very large classes of spaces the answer to this question, and some related questions, is ‘yes’. This can be thought of as a ‘ C^* -rigidity result’: it shows that the Roe algebra construction preserves a large amount of information about the space, and is thus surprisingly ‘rigid’.

As an example of our results, in the group case we have that if Γ and Λ are finitely generated elementary amenable, hyperbolic, or linear, groups such that the crossed products $l^\infty(\Gamma) \rtimes_r \Gamma$ and $l^\infty(\Lambda) \rtimes_r \Lambda$ are isomorphic, then Γ and Λ are quasi-isometric.

MSC: primary 46L85, 51K05.

Keywords: coarse geometry, coarse Baum-Connes conjecture.

1 Introduction

This piece asks to what extent certain noncommutative C^* -algebras associated to a metric space X , the *Roe-type algebras of X* , capture the large scale geometry of X . As such it forms part of the noncommutative geometry program of Connes [8]. It is also a contribution to the understanding of the *coarse Baum-Connes conjecture*, which predicts that, at least on the level of algebraic topology, Roe-type algebras are good models for the large scale geometry of spaces.

For us, ‘large scale geometry’ will mean those aspects of metric geometry that are preserved under coarse equivalence as in the following definition.

Definition 1.1. Let X and Y be metric spaces. A (not necessarily continuous) map $f : X \rightarrow Y$ is said to be *uniformly expansive* if for all $R > 0$ there exists $S > 0$ such that if $x_1, x_2 \in X$ satisfy $d(x_1, x_2) \leq R$, then $d(f(x_1), f(x_2)) \leq S$.

Two maps $f, g : X \rightarrow Y$ are said to be *close* if there exists $C > 0$ such that $d(f(x), g(x)) \leq C$ for all $x \in X$.

X and Y are said to be *coarsely equivalent* if there exist uniformly expansive maps $f : X \rightarrow Y$ and $g : Y \rightarrow X$ such that $f \circ g$ and $g \circ f$ are close to the identity maps on Y and X respectively.

Many metric spaces that are of interest from a classical point of view, for example finitely generated discrete groups with a word metric, or Riemannian manifolds of bounded sectional curvature and positive injectivity radius, are coarsely equivalent to uniformly discrete, bounded geometry metric spaces as in the following definition; in large scale geometry, it thus often suffices to consider metric spaces of this form.

Definition 1.2. Let X be a metric space. X is said to be *uniformly discrete* if there exists $\delta > 0$ such that if x, y are distinct points in X , then $d(x, y) \geq \delta$.

X is said to have *bounded geometry* if for all $R \geq 0$ there exists $N_R \in \mathbb{N}$ such that for all $x \in X$, the ball of radius R about x has at most N_R elements.

Throughout the remainder of this piece, the word *space* will mean uniformly discrete, bounded geometry metric space.

The *Roe-type algebras* that form our main object of study are defined below.

Definition 1.1. Let X be a space and $\mathcal{H}$ a separable (possibly finite dimensional) Hilbert space. Let $l^2(X, \mathcal{H})$ be the Hilbert space of square-summable functions from X to $\mathcal{H}$. Let T be a bounded operator on $l^2(X, \mathcal{H})$, which can be written as an X -by- X matrix

$$T = (T_{xy})_{x, y \in X}$$

where each T_{xy} is a bounded operator on $\mathcal{H}$.

A bounded operator T on $l^2(X, \mathcal{H})$ is called a *Roe operator* if the following two conditions hold.


- T has *finite propagation*: there exists $S > 0$ such that if $d(x, y) \geq S$ then $T_{xy} = 0$.
- All the matrix coefficients T_{xy} are rank one operators on $\mathcal{H}$.

A $*$ -subalgebra of $\mathcal{B}(l^2(X, \mathcal{H}))$ is called an *algebraic Roe-type algebra* if it consists only of finite propagation operators and contains all of the Roe operators. We denote algebraic Roe-type algebras by $A[X]$ (note however that if $\mathcal{H}$ is infinite dimensional, there are many non-isomorphic algebraic Roe-type algebras associated to X).

A C^* -subalgebra of $\mathcal{B}(l^2(X, \mathcal{H}))$ is called a *Roe-type algebra* if it is the norm closure of some algebraic Roe-type algebra. We denote Roe-type algebras by $A^*(X)$.

Let $\mathcal{H}$ be infinite dimensional. The *Roe algebra* of X is the completion of the $*$ -algebra of all finite propagation bounded operators T on $l^2(X, \mathcal{H})$ such that each matrix coefficient T_{xy} is a compact operator on $\mathcal{H}$.

The schematic below represents a Roe operator as an X -by- X matrix.



As we hope is clear from the picture, Roe operators can only ‘move’ elements of a space X a bounded distance, and are thus tied to its geometry.

Roe-type algebras were introduced by John Roe for the purpose of studying index theory on non-compact manifolds. See [25, 26] for the original articles, and [27, Chapter 3] and [28, Chapter 4] for more recent treatments. Roe-type algebras have since been extensively studied, both in their own right, and as they have applications to geometry and topology.

Many of these applications come via the *coarse Baum Connes conjecture*: the original sources are the articles of Roe [26, Conjecture 6.30], Higson–Roe [19], and Yu [36]. The coarse Baum-Connes conjecture has been very widely studied, and is responsible for some of the best results on problems such as the Novikov conjecture, and the existence of positive scalar curvature metrics.

Let X be a space, and $C^*(X)$ the Roe algebra of X . The coarse Baum-Connes conjecture asks whether a certain *assembly map*

$$\mu : \lim_{R \rightarrow \infty} K_*(P_R(X)) \rightarrow K_*(C^*(X))$$

is an isomorphism. For the present discussion, the left hand side should be thought of as the ‘large-scale algebraic topology of X ’ and the right hand side as the ‘algebraic topology of $C^*(X)$ ’. The conjecture asks whether certain natural geometric invariants (to be thought of as coming from elliptic differential operators on X) living in the *left hand side* are preserved by passage to the *right hand side*, where they have much better properties (e.g. vanishing properties, and homotopy invariance). In other words, the coarse Baum-Connes conjecture asks the following question.

Question 1.3. *If X and Y are metric spaces, how well does the algebraic topology of $C^*(X)$ model the large scale algebraic topology of X ?*

In this sense, the coarse Baum-Connes conjecture fits very naturally into the *noncommutative geometry* program of Alain Connes [8]: it postulates that the

noncommutative C^* -algebra $C^*(X)$ retains a significant amount of information about the space X , at least at the level of algebraic topology.

Since the formulation of the coarse Baum-Connes conjecture, it has become apparent that Roe-type algebras in general encode rather a lot of information about the large scale geometry of the underlying spaces: a particularly nice example is the theorem of Brown-Ozawa [4, Theorem 5.5.7] relating *nuclearity* of Roe-type algebras to *property A* for spaces (see also Guentner-Kaminker [14] and Ozawa [23]). Another very interesting example is given by results of Chen and Wang [6] that relate the ideal structure of the uniform Roe algebra $C_u^*(X)$ to the coarse geometric ‘ideal structure’ of X . See also [7, 5, 30, 31, 35] from this point of view.

All of this motivated the the following *rigidity question*. It can be thought of as a more precise analogue of Question 1.3 above: it asks about coarse geometric information at the level of algebras and spaces themselves, rather than merely at the level of algebraic-topological invariants associated to these algebras and spaces. A version of this question was also asked by Winter and Zacharias [35, page 3].

Question 1.4. *If X and Y are spaces which admit $*$ -isomorphic Roe-type algebras $A^*(X)$ and $A^*(Y)$, are X and Y coarsely equivalent?*

Question 1.4 is called a ‘rigidity question’ as it asks if the C^* -algebra $C^*(X)$ actually encodes *all* of the large scale geometry of X , and is thus surprisingly ‘rigid’. The terminology ‘rigid’ in this sort of context perhaps originates in the work of Mostow and Margulis on lattices in semisimple Lie groups.

The rigidity terminology is also motivated by an analogy between our question in coarse geometry / C^* -algebras and recent deep results of Popa and others (see [24] for a survey) in the theory of group actions / von Neumann algebras. One of the questions these authors consider is as follows: if the group-measure-space von Neumann algebras $L^\infty([0, 1]^\Gamma) \overline{\rtimes} \Gamma$ and $L^\infty([0, 1]^\Lambda) \overline{\rtimes} \Lambda$ associated to the Bernoulli shift actions of discrete groups Γ, Λ are isomorphic, then are the actions conjugate (and in particular, are the groups isomorphic)? If X is the coarse space underlying a discrete group Γ , then the ‘simplest’ Roe-type algebra associated to X (precisely, the uniform Roe algebra $C_u^*(X)$ – cf. Corollary 1.9 below) is isomorphic to the C^* -algebra crossed product $l^\infty(\Gamma) \rtimes_r \Gamma$, and is thus an analogue of the group-measure-space von Neumann algebra¹.

Our main results are that Question 1.4 and some variants often have a positive answer. To state the results precisely, we will need two properties of metric spaces.

The first is the *metric sparsification* property of Chen–Tessera–Wang–Yu [5, Section 3], as defined below; see Remarks 1.7 below for some comments.

¹The very recent results of Ioana [22] on *superrigidity* of Bernoulli shift actions have no obvious analogue in our context. Nonetheless, Thomas Sinclair has pointed out to us that there are some interesting similarities between our ideas and Ioana’s, despite the two being developed independently (in our context, the methods are also somewhat less analytically involved).

Definition 1.5. Let X be a space. X is said to have the *metric sparsification property* if there exists a constant $1 \geq \kappa > 0$ such that for all $S \geq 0$ there exists $D \geq 0$ such that for every Borel probability measure μ on X there exists a subset $\Omega \subseteq X$ equipped with a decomposition $\Omega = \sqcup_{i \in I} \Omega_i$ such that

- $\mu(\Omega) \geq \kappa$;
- $\text{diam}(\Omega_i) \leq D$ for all i ;
- $d(\Omega_i, \Omega_j) > S$ for all $i \neq j$.

The second property we need is Yu's *property A* [38, Section 2].

Definition 1.6. Let X be a space. X is said to have *property A* if for all $R, \epsilon > 0$ there exists a family of subsets $\{A_x\}_{x \in X}$ of $X \times \mathbb{N}$ such that:

- there exists $S \geq 0$ such that if $(x, n) \in A_y$, then $d(x, y) \leq S$;
- for all $x, y \in X$ such that $d(x, y) \leq R$, if $A_x \Delta A_y$ denotes the symmetric difference of A_x and A_y , then

$$\frac{|A_x \Delta A_y|}{|A_x \cap A_y|} < \epsilon.$$

Remarks 1.7. • Property A has been very widely studied (see [34] for a survey), while metric sparsification is a relatively new property [5, 16].

- Property A and the metric sparsification property are both implied by Gromov's *finite asymptotic dimension* [11, Section 1.E] – see [20, Lemma 4.3] and [5, Remark 3.2] respectively – and more generally by the Guentner–Tessera–Yu notion of *finite decomposition complexity* [15]. There are many examples of spaces with the metric sparsification property, and property A, that do not have finite asymptotic dimension, however: a simple example is the wreath product $\mathbb{Z} \wr \mathbb{Z}$ equipped with a word metric.
- Property A and metric sparsification both hold for a very large class of spaces: notable examples in the group case are hyperbolic groups [29], elementary amenable groups, and linear groups [13, 16].
- The metric sparsification property and property A both fail for a coarse disjoint union of expanding graphs with uniform expansion constant and cardinality tending to infinity. Relatedly, the only *groups* for which these properties are known to fail are the random groups of Gromov whose Cayley graphs coarsely contain such a sequence of expanding graphs [12, 1].
- The precise relationship between the metric sparsification property and property A is not known: it seems an interesting open problem to clarify this. One possible direction here is that property A is known for all amenable groups, but metric sparsification is only known for elementary amenable groups.

Our main results can then be summarised as follows.

Theorem 1.8. *Let X and Y be spaces.*

1. *If X and Y have the metric sparsification property and admit $*$ -isomorphic Roe-type algebras $A^*(X)$ and $A^*(Y)$, then X and Y are coarsely equivalent.*
2. *If X and Y have property A and admit $*$ -isomorphic algebraic Roe-type algebras $A[X]$ and $A[Y]$, then X and Y are coarsely equivalent.*
3. *If X and Y admit Roe-type algebras $A[X]$ and $A[Y]$ that are $*$ -isomorphic via a map that extends to a $*$ -isomorphism of $A^*(X)$ with $A^*(Y)$ (equivalently, that is spatially implemented), then X and Y are coarsely equivalent.*

As an interesting corollary, we note the following result.

Corollary 1.9. *Say Γ and Λ are countable (respectively, finitely generated) groups which have the metric sparsification property when equipped with some (equivalently, any) proper left-invariant metric. If $l^\infty(\Gamma) \rtimes_r \Gamma \cong l^\infty(\Lambda) \rtimes_r \Lambda$, then Γ and Λ are coarsely equivalent (respectively, quasi-isometric).*

Proof. For Γ a countable discrete group, the crossed product $l^\infty(\Gamma) \rtimes_r \Gamma$ is an algebra of Roe-type: it is in fact isomorphic to the uniform Roe algebra $C_u^*(|\Gamma|)$ (this observation is due to Higson and Yu, see for example [4, Proposition 5.1.3]). Part 1 of Theorem 1.8 then implies that Γ and Λ are coarsely equivalent; moreover, for finitely generated groups, coarse equivalence is the same as quasi-isometry. $\square$

These results follow from Theorems 4.1, 5.1, 5.2 and 6.1 below.

We conclude this introduction with some remarks on Theorem 1.8.

Remarks 1.10. • Part 2 has a purely algebraic statement (particularly if $\mathcal{H} = \mathbb{C}$) and could be thought of in terms of the *translation algebras* used in controlled topology and elsewhere (see [28, Chapter 4] for an introduction). Nonetheless, the proof we give makes essential use of analytic and operator algebraic ideas.

- There are several natural Roe-type algebras that have been introduced for the study of index theory and coarse geometry: see Examples 2.1 and Example 2.2 below. Theorem 1.8 is of most interest in these special cases; we have introduced the framework of Roe-type algebras mainly to make the assumptions clearer, and to streamline the exposition.
- It follows from our methods that, up to a natural notion of equivalence in each case, coarse equivalences are essentially the same thing as spatially implemented isomorphisms of algebraic Roe algebras. This is made precise in Appendix A.

- The coarse Baum-Connes conjecture is false in general: see [37, Section 8] and [17, 18]. Our work here is a step towards a more precise geometric understanding of why one should expect it to be true in ‘good’ cases. We hope to be able to expand on this project in order to elucidate what goes wrong in more exotic cases. A promising direction here seems to be provided by the *ghost ideals* studied in [6, 7, 33] and [28, Section 11.5.2].

Outline of the piece

Section 2 gives definitions, recalls some standard notation that we need, and gives the main examples of Roe-type algebras. Section 3 covers some analytic preliminaries: the first is an easy sufficient condition for a $*$ -isomorphism to be spatially implemented (this, however, is crucial to get the machinery going in the first place); the second shows that ‘cancellation of arbitrarily large propagation’ cannot occur when one sums operators in a Roe algebra (this is perhaps the main ingredient in our results). Section 4 proves part 1 of Theorem 1.8, and Section 5 proves parts 2 and 3. There are some slight additional subtleties that mean that the *stable uniform Roe algebra* $C_s^*(X)$ does not fit into our general framework; the adjustments needed to apply our methods to it are dealt with in Section 6. Finally, Appendix A defines categories of spaces in which the morphisms are (equivalence classes of) coarse equivalences and (equivalence classes of) spatially implemented isomorphisms of Roe algebras, and shows that these two categories are isomorphic.

2 Definitions and notation

We start this section by looking more precisely at our definition of Roe-type algebras (Definition 1.1 from the introduction).

We first look at our conventions on Hilbert spaces. All Hilbert spaces considered in this piece are separable and complex, and inner products are linear in the second variable, conjugate linear in the first. If $\mathcal{H}$ is a Hilbert space, we write

$$\mathcal{H}_1 := \{v \in \mathcal{H} \mid \|v\| \leq 1\}$$

for the closed unit ball in $\mathcal{H}$. We write $\mathcal{B}(\mathcal{H})$ (respectively, $\mathcal{K}(\mathcal{H})$) for the C^* -algebra of bounded (respectively, compact) operators on $\mathcal{H}$. If X is a set and $\mathcal{H}$ a Hilbert space, $l^2(X, \mathcal{H})$ denotes the Hilbert space of square summable functions on X with values in $\mathcal{H}$. $l^2(X)$ denotes $l^2(X, \mathbb{C})$, and the standard orthonormal basis of this space is denoted $\{\delta_x\}_{x \in X}$. If $l^2(X) \otimes \mathcal{H}$ is the (completed) Hilbert space tensor product, then there is a natural isomorphism

$$l^2(X) \otimes \mathcal{H} \cong l^2(X, \mathcal{H})$$

defined by sending the elementary tensor $\delta_x \otimes v$ to the function from X to $\mathcal{H}$ defined by

$$y \mapsto \begin{cases} v & y = x \\ 0 & \text{otherwise} \end{cases} ;$$

we will often make use of this isomorphism, and the corresponding natural representation of the spatial tensor product $\mathcal{B}(l^2(X)) \otimes \mathcal{B}(\mathcal{H})$ on $l^2(X, \mathcal{H})$, without further comment.

If A is a subset of X we write χ_A for the characteristic function of A considered as a multiplication operator on $l^2(X, \mathcal{H})$. For $x, y \in X$ we write e_{xy} for the ‘matrix unit’ in $\mathcal{B}(l^2(X, \mathcal{H}))$ defined on elementary tensors by

$$e_{xy} : \delta_z \otimes v \mapsto \langle \delta_y, \delta_z \rangle \delta_x \otimes v.$$

Similarly, if $x, y \in X$ and $v, w \in \mathcal{H}$, we write $e_{(x,v),(y,w)} \in \mathcal{B}(l^2(X, \mathcal{H}))$ for the rank one ‘matrix unit’ defined by

$$e_{(x,v),(y,w)} : \delta_z \otimes u \mapsto \langle \delta_y \otimes w, \delta_z \otimes u \rangle \delta_x \otimes v. \quad (1)$$

If T is a bounded operator from $l^2(Y, \mathcal{H})$ to $l^2(X, \mathcal{H})$, $x \in X$ and $y \in Y$, we write

$$T_{xy} := e_{xx} T e_{yy},$$

for the ‘ $(x, y)^{\text{th}}$ matrix coefficient of T ’, which naturally identifies with a bounded operator on $\mathcal{H}$. Similarly if $x \in X$, $y \in Y$ and $v, w \in \mathcal{H}$, we write

$$T_{(x,v),(y,w)} := \langle \delta_x \otimes v, T(\delta_y \otimes w) \rangle \quad (2)$$

for the ‘ $((x, v), (y, w))^{\text{th}}$ matrix coefficient’ of T .

The following gives some examples of Roe-type algebras as in Definition 1.1; all have applications in coarse geometry and index theory.

Examples 2.1. Let X be a space and $\mathcal{H}$ be a separable Hilbert space. The following $*$ -algebras of bounded operators on $l^2(X, \mathcal{H})$ are all algebraic Roe-type algebras:

- say $\mathcal{H} = \mathbb{C}$ and let $\mathbb{C}_u[X]$ consist of all finite propagation operators on $l^2(X, \mathcal{H}) = l^2(X)$;
- say $\mathcal{H}$ is infinite dimensional and let $UC[X]$ consist of all finite propagation operators T such that there exists $N \in \mathbb{N}$ such that for all $x, y \in X$, T_{xy} is an operator of rank at most N ;
- say $\mathcal{H}$ is infinite dimensional and let $\mathbb{C}[X]$ consist of finite propagation operators T such that for all $x, y \in X$, T_{xy} is in $\mathcal{K}(\mathcal{H})$.

The operator norm completions of these three $*$ -algebras are denoted $C_u^*(X)$, $UC^*(X)$ and $C^*(X)$ respectively, and are called the *uniform Roe algebra of X* , *uniform algebra of X* , and *Roe algebra of X* , respectively. They are, of course, all Roe-type algebras.

The following example does not quite fit into the framework above. However, our methods apply to it with some minor modifications (see Section 6 below); in particular, the obvious analogue of Theorem 1.8 is true in this case.

Example 2.2. Let X be a space and $\mathcal{H}$ an infinite dimensional Hilbert space. $\mathbb{C}_s[X]$ consists of all finite propagation operators T on $l^2(X, \mathcal{H})$ such that there exists a finite dimensional subspace $\mathcal{H}_T \subseteq \mathcal{H}$ such that for all $x, y \in X$, $T_{xy} \in \mathcal{B}(\mathcal{H}_T)$. The operator norm closure of $\mathbb{C}_s[X]$ is denoted $C_s^*(X)$ and called the *stable uniform Roe algebra* of X .

Remarks 2.3. 1. It is not hard to see that $C_s^*(X)$ is canonically isomorphic to $C_u^*(X) \otimes \mathcal{K}(\mathcal{H})$, whence its name.

2. Clearly one has inclusions

$$C_u^*(X) \subseteq C_s^*(X) \subseteq UC^*(X) \subseteq C^*(X),$$

the last three of which are canonical. In general, none of these inclusions are equalities, and moreover none of these four C^* -algebras associated to X is abstractly isomorphic to any of the others.

3. It will be important in what follows to note that any algebraic Roe-type algebra associated to a space X contains all the rank one operators $e_{(x,v),(y,w)}$, where $x, y \in X$ and $v, w \in \mathcal{H}$. It follows in particular that any Roe-type algebra associated to X contains the compact operators $\mathcal{K}(l^2(X, \mathcal{H}))$ as a C^* -subalgebra. The same remarks apply to $\mathbb{C}_s[X]$ and $C_s^*(X)$: the former contains all the rank-one operators $e_{(x,v),(y,w)}$, and the latter contains the compact operators.
4. The Roe-type algebras $C_s^*(X)$, $UC^*(X)$ and $C^*(X)$ are all coarse invariants, in the sense that if X and Y are coarsely equivalent spaces, then $C_s^*(X) \cong C_s^*(Y)$, $UC^*(X) \cong UC^*(Y)$ and $C^*(X) \cong C^*(Y)$. Similar statements hold on the purely algebraic level, but not for $C_u^*(X)$ as one can see by considering finite spaces. In these three cases, the various parts of Theorem 1.8 above thus have converses: this is folklore for $C^*(X)$ and $UC^*(X)$, and follows from [2, Theorem 4] for $C_s^*(X)$.
5. The Roe-type algebras $UC^*(X)$ and $C^*(X)$, and the stable uniform Roe algebra $C_s^*(X)$, can be made sense of for metric spaces that are coarsely equivalent to bounded geometry, uniformly discrete spaces, rather than metric spaces that actually have these properties themselves. Theorem 1.8 still holds in this context.

3 Analytic Preliminaries

The following lemma is a version of well-known classical ideas. We could not find a reference for exactly what we needed in the literature, however, so provide a proof.

Lemma 3.1. *Let A and B be C^* -algebras, concretely represented on Hilbert spaces $\mathcal{H}_A$ and $\mathcal{H}_B$ respectively, and such that A (resp. B) contains the compact operators $\mathcal{K}(\mathcal{H}_A)$ (resp. $\mathcal{K}(\mathcal{H}_B)$).*

Then any $*$ -isomorphism $\phi : A \rightarrow B$ is spatially implemented, i.e. there exists a unitary isomorphism $U : \mathcal{H}_A \rightarrow \mathcal{H}_B$ such that

$$\phi(T) = UTU^*$$

for all $T \in A$.

Proof. Note first that as $\mathcal{K}(\mathcal{H}_A)$ is an essential, minimal ideal in $\mathcal{B}(\mathcal{H}_A)$, $\mathcal{K}(\mathcal{H}_A)$ must be the unique minimal ideal in A , and similarly for B . Hence ϕ must restrict to a $*$ -isomorphism from $\mathcal{K}(\mathcal{H}_A)$ onto $\mathcal{K}(\mathcal{H}_B)$. Any such $*$ -isomorphism is spatially implemented (see for example [9, Corollary 4.1.8]), whence there exists a unitary isomorphism $U : \mathcal{H}_A \rightarrow \mathcal{H}_B$ such that

$$\phi(K) = UKU^*$$

for all $K \in \mathcal{K}(\mathcal{H}_A)$.

Recall now that for a sequence (T_n) of bounded operators on $\mathcal{H}_A$, T_n converges to some T in the strong operator topology if and only if $T_n K$ converges to TK in the norm topology for all compact operators K . Hence if (T_n) is a sequence in A converging to some $T \in A$ in the strong operator topology then, as ϕ is norm continuous and restricts to an isomorphism $\mathcal{K}(\mathcal{H}_A) \rightarrow \mathcal{K}(\mathcal{H}_B)$, we must have that $\phi(T_n) \rightarrow \phi(T)$ in the strong operator topology on $\mathcal{B}(\mathcal{H}_B)$.

Finally, write any $T \in A$ as a strong limit of a sequence T_n of compact operators (which one can do using cut-downs by finite rank projections, for example). We have that

$$\phi(T) = \lim_n \phi(T_n) = \lim_n UT_n U^* = U(\lim_n T_n)U^* = UTU^*$$

as required. $\square$

The next lemma, and the similar Lemma 3.3, are the most important tools in proving our main results.

Lemma 3.2. *Let X be a space, and $A^*(X)$ denote either a Roe-type algebra associated to X , or the stable uniform Roe algebra $C_s^*(X)$.*

Let $(T_n)_{n \in \mathbb{N}}$ be a sequence of mutually orthogonal operators in $A^*(X)$, and assume that for any $\mathcal{N} \subseteq \mathbb{N}$, the sum $\sum_{n \in \mathcal{N}} T_n$ converges strongly to some operator in $A^*(X)$.

Then for any $\epsilon > 0$ there exists $S \geq 0$ such that for any subsets A, B of X with $d(A, B) > S$, and any n , we have that

$$\|\chi_A T_n \chi_B\| < \epsilon.$$

Proof. Choose a sequence (p_k) of finite rank projections in $\mathcal{H}$ that converge strongly to the identity, and let

$$P_k = 1 \otimes p_k \in l^2(X) \otimes \mathcal{H}.$$

Choose also sequences of nested finite subsets $(A_k), (B_k)$ of A, B respectively such that $A = \cup A_k$ and $B = \cup B_k$. As for any n we have that

$$(\chi_{A_k} P_k) T_n (\chi_{B_k} P_k) \xrightarrow{s.o.t.} \chi_A T_n \chi_B \quad \text{as } k \rightarrow \infty$$

(this uses that the sequences $(\chi_{A_k} P_k)$ and $(\chi_{B_k} P_k)$ are uniformly bounded), and as limits in the strong operator topology do not increase norms, it suffices to prove the following statement: ‘for any $\epsilon > 0$ there exists $S \geq 0$ such that for any finite subsets A, B of X such that $d(A, B) > S$, all k and any n , we have that

$$\|\chi_A P_k T_n \chi_B P_k\| < \epsilon.$$

We will prove this statement.

Assume for contradiction that there is $\epsilon > 0$, such that for any $S \geq 0$ there are S -separated finite subsets $A_S, B_S \subseteq X$, a projection $P_{k_S} = 1 \otimes p_{k_S}$ and i_S , such that

$$\|\chi_{A_S} P_{k_S} T_{i_S} \chi_{B_S} P_{k_S}\| \geq \epsilon.$$

Choose any sequence (S_n) tending to infinity, and for simplicity rename $A_{S_n}, B_{S_n}, P_{k_{S_n}}$ and i_{S_n} as A_n, B_n, P_n and n respectively.

We will show that there are two subsets $\mathcal{N}, \mathcal{M} \subseteq \mathbb{N}$ such that:

- $\lim_{n \in \mathcal{N}} S_n = \infty$ (in other words, $\mathcal{N}$ is infinite); and
- for all $n \in \mathcal{N}$,

$$\left\| \chi_{A_n} P_n \left(\sum_{m \in \mathcal{M}} T_m \right) \chi_{B_n} P_n \right\| \geq \frac{\epsilon}{4}.$$

This contradicts the fact that $\sum_{m \in \mathcal{M}} T_m$ is in $A^*(X)$.

We define $\mathcal{N}$ and $\mathcal{M}$ inductively; start with $\mathcal{N} = \mathcal{M} = \emptyset$, and a parameter $k = 1$. For the inductive step, we assume that we have ‘updated’ $\mathcal{N}, \mathcal{M}$ and k . We declare the updated $\mathcal{N}$ to be $\mathcal{N} \cup \{k\}$. Now, by assumption,

$$\|\chi_{A_k} P_k T_k \chi_{B_k} P_k\| \geq \epsilon,$$

whence either $\mathcal{M}' := \mathcal{M}$ or $\mathcal{M}' := \mathcal{M} \cup \{k\}$ satisfies

$$\left\| \chi_{A_k} P_k \left(\sum_{m \in \mathcal{M}'} T_m \right) \chi_{B_k} P_k \right\| \geq \frac{\epsilon}{2}.$$

We rename $\mathcal{M}'$ to $\mathcal{M}$. As the operators T_m are mutually orthogonal and the sum $\sum_{m > k} T_m$ converges strongly, the sum

$$\sum_{m > k} \chi_{A_k} P_k T_{\sigma(m)} \chi_{B_k} P_k \tag{3}$$

converges strongly for any permutation σ of $\{m \in \mathbb{N} \mid m > k\}$; the Riemann rearrangement theorem for the finite dimensional vector space

$$\mathcal{B}(\chi_{B_k} P_k \cdot (l^2(X, \mathcal{H})), \chi_{A_k} P_k \cdot (l^2(X, \mathcal{H})))$$

thus implies that the sum in line (3) above converges absolutely; in particular, there exists $k' > k$ such that

$$\sum_{m \geq k'} \|\chi_{A_k} P_k T_m \chi_{B_k} P_k\| \leq \frac{\epsilon}{4}.$$

In particular, if $\mathcal{M}'$ is the result of adding any number of elements to $\mathcal{M}$, all of which are larger than k' , then

$$\left\| \chi_{A_k} P_k \left(\sum_{m \in \mathcal{M}'} T_m \right) \chi_{B_k} P_k \right\| \geq \frac{\epsilon}{2} - \frac{\epsilon}{4} = \frac{\epsilon}{4}.$$

We declare the new k to be k' and repeat the induction. It is clear that this process yields the required subsets $\mathcal{N}$ and $\mathcal{M}$. $\square$

We will also need the following slightly different version of the above lemma; as the proof is essentially the same, we omit it.

Lemma 3.3. *Let X be a space, and let $A[X]$ either denote an algebraic Roe-type algebra associated to X , or let $A[X]$ denote $\mathbb{C}_s[X]$.*

Let $(T_n)_{n \in \mathbb{N}}$ be a sequence of mutually orthogonal operators in $A[X]$, and assume that for any $\mathcal{N} \subseteq \mathbb{N}$, the sum $\sum_{n \in \mathcal{N}} T_n$ converges strongly to some operator in $A[X]$.

Then there exists $S \geq 0$ such that for any subsets A, B of X with $d(A, B) > S$, and any n , we have that

$$\chi_A T_n \chi_B = 0. \quad \square$$

4 C^* -algebraic isomorphisms and coarse equivalences

In this section, we will prove the following theorem.

Theorem 4.1. *Say X and Y are spaces with the metric sparsification property, and $A^*(X)$ and $A^*(Y)$ are Roe-type algebras associated to X and Y respectively.*

If there exists a $*$ -isomorphism $\phi : A^*(X) \rightarrow A^*(Y)$, then X and Y are coarsely equivalent.

For the remainder of this section, fix X, Y and $\phi : A^*(X) \rightarrow A^*(Y)$ as in the statement of Theorem 4.1. Note that Lemma 3.1 (together with part 3 of Remark 2.3) implies that there exists a unitary isomorphism $U : l^2(X, \mathcal{H}) \rightarrow l^2(Y, \mathcal{H})$ spatially implementing ϕ ; we also fix U throughout.

The following very simple lemma is isolated for ease of reference; see lines (1) and (2) above for notation.

Lemma 4.2. *For any $x_1, x_2 \in X$, $v_1, v_2 \in \mathcal{H}$, $y_1, y_2 \in Y$, and $w_1, w_2 \in \mathcal{H}$ we have the formula*

$$(\phi(e_{(x_1, v_1), (x_2, v_2)}))_{(y_1, w_1), (y_2, w_2)} = \langle \delta_{y_1} \otimes w_1, U(\delta_{x_1} \otimes v_1) \rangle \langle U(\delta_{x_2} \otimes v_2), \delta_{y_2} \otimes w_2 \rangle.$$

Proof. Computing,

$$\begin{aligned}
& (\phi(e_{(x_1, v_1), (x_2, v_2)}))_{(y_1, w_1), (y_2, w_2)} \\
&= \langle \delta_{y_1} \otimes w_1, \phi(e_{(x_1, v_1), (x_2, v_2)})(\delta_{y_2} \otimes w_2) \rangle \\
&= \langle U^*(\delta_{y_1} \otimes w_1), e_{(x_1, v_1), (x_2, v_2)} U^*(\delta_{y_2} \otimes w_2) \rangle \\
&= \langle U^*(\delta_{y_1} \otimes w_1), \langle \delta_{x_2} \otimes v_2, U^*(\delta_{y_2} \otimes w_2) \rangle \delta_{x_1} \otimes v_1 \rangle \\
&= \langle \delta_{x_2} \otimes v_2, U^*(\delta_{y_2} \otimes w_2) \rangle \langle U^*(\delta_{y_1} \otimes w_1), \delta_{x_1} \otimes v_1 \rangle \\
&= \langle U(\delta_{x_2} \otimes v_2), \delta_{y_2} \otimes w_2 \rangle \langle \delta_{y_1} \otimes w_1, U(\delta_{x_1} \otimes v_1) \rangle
\end{aligned}$$

as required. $\square$

Lemma 4.3. 1. Let $\delta > 0$. Then there exists $S \geq 0$ such that if $x \in X$, $v_1, v_2 \in \mathcal{H}_1$, $y_1, y_2 \in Y$ and $w_1, w_2 \in \mathcal{H}_1$ are such that

$$|\langle U(\delta_x \otimes v_1), \delta_{y_1} \otimes w_1 \rangle| \geq \delta \quad \text{and} \quad |\langle U(\delta_x \otimes v_2), \delta_{y_2} \otimes w_2 \rangle| \geq \delta,$$

then $d(y_1, y_2) \leq S$.

2. Let $\delta > 0$. Then for all $R > 0$ there exists $S \geq 0$ such that if $x_1, x_2 \in X$ are such that $d(x_1, x_2) \leq R$ and $v_1, v_2 \in \mathcal{H}_1$, $y_1, y_2 \in Y$, and $w_1, w_2 \in \mathcal{H}_1$ are such that

$$|\langle U(\delta_{x_1} \otimes v_1), \delta_{y_1} \otimes w_1 \rangle| \geq \delta \quad \text{and} \quad |\langle U(\delta_{x_2} \otimes v_2), \delta_{y_2} \otimes w_2 \rangle| \geq \delta,$$

then $d(y_1, y_2) \leq S$.

The same properties hold with the roles of X and Y reversed, and with U replaced by U^* .

Proof. The fact that the same properties hold with the roles of X and Y interchanged follows by symmetry. Moreover, the first of these properties is a special case of the second. It thus suffices to prove the second property in the form stated.

Assume for contradiction that this is false. Then there exists $R \geq 0$ and sequences $(x_1^n), (x_2^n), (v_1^n), (v_2^n), (y_1^n), (y_2^n), (w_1^n), (w_2^n)$ such that $d(x_1^n, x_2^n) \leq R$ for all n , so that

$$|\langle U(\delta_{x_1^n} \otimes v_1^n), \delta_{y_1^n} \otimes w_1^n \rangle| \geq \delta \quad \text{and} \quad |\langle U(\delta_{x_2^n} \otimes v_2^n), \delta_{y_2^n} \otimes w_2^n \rangle| \geq \delta,$$

and so that $d(y_1^n, y_2^n) \rightarrow \infty$ as $n \rightarrow \infty$. Using bounded geometry of X , we may assume that the elements x_1^n, x_2^n are all distinct, whence the operators $e_{(x_1^n, v_1^n), (x_2^n, v_2^n)}$ are mutually orthogonal and for any $\mathcal{N} \subseteq \mathbb{N}$ the sum

$$\sum_{n \in \mathcal{N}} e_{(x_1^n, v_1^n), (x_2^n, v_2^n)}$$

converges strongly to an element in $A^*(X)^2$. Hence the same is true of the sums

$$\sum_{n \in \mathcal{N}} \phi(e_{(x_1^n, v_1^n), (x_2^n, v_2^n)}),$$

²The fact that this operator need not exist in $C_s^*(X)$ is the reason for treating that case separately.

in $A^*(Y)$, using strong continuity of ϕ (which follows from Lemma 3.1). Lemma 3.2 thus implies that there exists $S > 0$ such that for any $A, B \subseteq Y$ with $d(A, B) > S$ we have that

$$\|\chi_A \phi(e_{(x_1^n, v_1^n), (x_2^n, v_2^n)}) \chi_B\| < \delta^2.$$

Set $A_n = \{y_1^n\}$, $B_n = \{y_2^n\}$. Then we have that for all n so large that $d(y_1^n, y_2^n) > S$

$$\begin{aligned} \|\chi_{A_n} \phi(e_{(x_1^n, v_1^n), (x_2^n, v_2^n)}) \chi_{B_n}\| &\geq \|\phi(e_{(x_1^n, v_1^n), (x_2^n, v_2^n)})_{(y_1^n, w_1^n), (y_2^n, w_2^n)}\| \\ &= |\langle \delta_{y_1^n} \otimes w_1^n, U(\delta_{x_1^n} \otimes v_1^n) \rangle \langle U(\delta_{x_2^n} \otimes v_2^n), \delta_{y_2^n} \otimes w_2^n \rangle|, \end{aligned}$$

where the last equality uses Lemma 4.2. This is greater than δ^2 by assumption, however, which is a contradiction. $\square$

The following analytic lemma will be our only use of the metric sparsification property.

Lemma 4.4. *Fix a unit vector $v_0 \in \mathcal{H}$. Then there exists $c > 0$ such that for each $x \in X$ there exists $f(x) \in Y$ and $w_x \in \mathcal{H}_1$ such that*

$$|\langle U(\delta_x \otimes v_0), \delta_{f(x)} \otimes w_x \rangle| \geq c$$

Remark 4.5. In fact, c exists independently of the choice of v_0 , but the slightly weaker statement above applies verbatim to the case considered in Section 6, and is all we need.

Proof. Let κ be as in Definition 1.5. Let $\epsilon = \kappa/4$ and $t = \kappa/5$, so in particular we have that

$$t + \epsilon < \frac{\kappa}{2}. \quad (4)$$

Applying Lemma 3.2 to the sum

$$\sum_{x \in X} \phi(e_{(x, v_0), (x, v_0)}) \in A^*(X)$$

(similarly to the proof of Lemma 4.3 above) implies that there exists $S > 0$ so that for any $x \in X$ and $A, B \subseteq Y$ with $d(A, B) > S$ we have

$$\|\chi_A \phi(e_{(x, v_0), (x, v_0)}) \chi_B\| < \epsilon. \quad (5)$$

For the remainder of the proof, fix $x \in X$. Define $\xi = U(\delta_x \otimes v_0)$, which we think of as a function from Y to $\mathcal{H}$. Note that $\chi_A \phi(e_{(x, v_0), (x, v_0)}) \chi_B$ is equal to the rank one operator on $l^2(Y, \mathcal{H})$ defined by

$$\eta \mapsto \langle \xi|_B, \eta \rangle \xi|_A$$

(where ‘ $|_C$ ’ denotes restriction to a subset $C \subseteq Y$), whence line (5) above implies that

$$\|\xi|_A\| \|\xi|_B\| < \epsilon. \quad (6)$$

Now, using the metric sparsification property for X applied to the Borel measure (in the current context, just a non-negative function with total mass one)

$$\begin{aligned}\mu : X &\rightarrow [0, 1] \\ x &\mapsto \|\xi(x)\|^2\end{aligned}$$

there exists a subset $\Omega = \sqcup_{i \in I} \Omega_i$ of X and $D \geq 0$ such that $\text{diam}(\Omega_i) \leq D$ for all i , $d(\Omega_i, \Omega_j) > S$ for all $i \neq j$ and

$$\|\xi|_{\Omega}\|^2 \geq \kappa.$$

Assume for contradiction that $\|\xi|_{\Omega_i}\|^2 < t$ for all i . Then there exists a partition $I = I_1 \sqcup I_2$ such that

$$\sum_{i \in I_1} \|\xi|_{\Omega_i}\|^2 \geq \frac{\kappa}{2} - t \quad \text{and} \quad \sum_{i \in I_2} \|\xi|_{\Omega_i}\|^2 \geq \frac{\kappa}{2} - t.$$

Taking $A = \sqcup_{i \in I_1} \Omega_i$ and $B = \sqcup_{i \in I_2} \Omega_i$, line (6) above implies that

$$\frac{\kappa}{2} - t \leq \|\xi|_A\| \|\xi|_B\| < \epsilon,$$

which contradicts line (4) above.

Hence $\|\xi|_{\Omega_i}\|^2 \geq t$ for some i . Using bounded geometry and uniform boundedness of the sets Ω_i , there is some $M \in \mathbb{N}$ (that depends only on S) such that $|\Omega_i| \leq M$ for all i . Hence there is $y \in \Omega_i$ such that

$$\|\xi(y)\|^2 \geq t/M.$$

The lemma is then true for $f(x)$ equal to this y , $w_x = \xi(y)/\|\xi(y)\|$, and $c = \sqrt{t/M}$. $\square$

Proof of Theorem 4.1. Let v_0 denote some fixed choice of unit vector in $\mathcal{H}$, and let $c > 0$ have the property in Lemma 4.4 both as stated, and with the roles of X and Y reversed. Define a function $f : X \rightarrow Y$ by choosing $f(x)$ to be any element of Y such that there exists $w_x \in \mathcal{H}_1$ such that

$$|\langle U(\delta_x \otimes v_0), \delta_{f(x)} \otimes w_x \rangle| \geq c$$

(such exists by Lemma 4.4). Similarly, define a function $g : Y \rightarrow X$ by choosing $g(y)$ to be any element of X such that there exists $v_y \in \mathcal{H}_1$ such that

$$|\langle U^*(\delta_y \otimes v_0), \delta_{g(y)} \otimes v_y \rangle| \geq c.$$

Note now that Lemma 4.3, part 2, implies that both f and g are uniformly expansive. Moreover, for any $x \in X$, we have both

$$|\langle U^*(\delta_{f(x)} \otimes v_0), \delta_{g(f(x))} \otimes v_{f(x)} \rangle| \geq c \quad \text{and} \quad |\langle U(\delta_x \otimes v_0), \delta_{f(x)} \otimes w_x \rangle| \geq c,$$

whence Lemma 4.3, part 1, implies the existence of some S (independently of x) such that

$$d(x, g(f(x))) \leq S.$$

This says that $g \circ f$ is close to the identity. Similarly, $f \circ g$ is close to the identity, and we are done. $\square$

5 Algebraic isomorphisms and coarse equivalences

In this section, we will prove the following theorem.

Theorem 5.1. *Say X and Y are spaces with property A , and $A[X]$ and $A[Y]$ are algebraic Roe-type algebras associated to X and Y respectively.*

If there exists a $*$ -isomorphism $\phi : A[X] \rightarrow A[Y]$, then X and Y are coarsely equivalent.

We will not need to use property A itself, just a consequence of it from [32, Section 2]. In fact, the only time property A is used in the proof of the above theorem is Lemma 5.4 below: this shows that in the presence of property A a $*$ -isomorphism as in Theorem 5.1 above extends to a $*$ -isomorphism on the completions

$$\tilde{\phi} : A^*(X) \rightarrow A^*(Y);$$

Lemma 3.1 then implies that ϕ is spatially implemented by a unitary isomorphism $U : l^2(X, \mathcal{H}) \rightarrow l^2(Y, \mathcal{H})$. The methods in this section thus also prove the following result.

Theorem 5.2. *Say X and Y are spaces with property A , and $A[X]$ and $A[Y]$ are algebraic Roe-type algebras associated to X and Y respectively.*

If there exists a $*$ -isomorphism $\phi : A[X] \rightarrow A[Y]$ that is spatially implemented by a unitary isomorphism $U : l^2(X, \mathcal{H}) \rightarrow l^2(Y, \mathcal{H})$ (equivalently, that extends to a $*$ -isomorphism $\tilde{\phi} : A^*(X) \rightarrow A^*(Y)$), then X and Y are coarsely equivalent. $\square$

We will use the following definition (due to Gong, Wang, and Yu [10]) as a crutch to Theorem 5.1 above.

Definition 5.3. Let X be a space, and $A[X]$ an algebraic Roe-type algebra associated to X . The corresponding *maximal completion* of $A[X]$, denoted $A_{max}^*(X)$, is the completion of $A[X]$ for the norm

$$\|T\|_{max} := \sup\{\|\pi(T)\|_{\mathcal{B}(\mathcal{H}')} \mid \pi : A[X] \rightarrow \mathcal{B}(\mathcal{H}') \text{ a } * \text{-homomorphism}\}$$

(contrary to our usual conventions, we allow the Hilbert spaces $\mathcal{H}'$ in the above to have uncountable dimension).

See [10, Lemma 3.4] for a proof that this norm is finite – that proof is only for the case of $\mathbb{C}[X]$, but the same idea works in general.

Lemma 5.4. *Say X and Y are spaces with property A , and $A[X]$ and $A[Y]$ are algebraic Roe-type algebras associated to X and Y respectively. Let $\phi : A[X] \rightarrow A[Y]$ be a $*$ -isomorphism. Let $A^*(X)$ and $A^*(Y)$ be the C^* -algebraic completions of $A[X]$ and $A[Y]$ respectively. Then ϕ extends (uniquely) to a $*$ -isomorphism $\tilde{\phi} : A^*(X) \rightarrow A^*(Y)$.*

Proof. Note first that if Z is a space, $A[Z]$ an associated algebraic Roe-type algebra, B any C^* -algebra, and $\psi : A[Z] \rightarrow B$ a $*$ -homomorphism, then ψ extends (uniquely) to a $*$ -homomorphism $\tilde{\psi} : A_{max}^*(Z) \rightarrow B$. It follows that ϕ as in the statement extends to a $*$ -isomorphism

$$\tilde{\phi} : A_{max}^*(X) \xrightarrow{\cong} A_{max}^*(Y). \quad (7)$$

Property A for a space Z , however, implies that the canonical quotient map

$$A_{max}^*(Z) \rightarrow A^*(Z)$$

is an isomorphism: see [32, Proposition 1.3] (this only covers the $C^*(Z)$ and $C_u^*(Z)$ cases, but the same proof works in general). Hence line (7) above identifies naturally with a $*$ -isomorphism

$$\tilde{\phi} : A^*(X) \xrightarrow{\cong} A^*(Y)$$

extending ϕ . □

It follows from Lemma 3.1 that $\tilde{\phi}$ is spatially implemented by a unitary isomorphism $U : l^2(X, \mathcal{H}) \rightarrow l^2(Y, \mathcal{H})$; we fix U for the rest of this section.

The following lemma is an analogue of Lemma 4.3. The proof is essentially the same (using Lemmas 3.3 and 4.2), so we omit it.

Lemma 5.5. *1. There exists $S \geq 0$ such that if $x \in X$, $v_1, v_2 \in \mathcal{H}_1$, $y_1, y_2 \in Y$ and $w_1, w_2 \in \mathcal{H}_1$ are such that*

$$|\langle U(\delta_x \otimes v_1), \delta_{y_1} \otimes w_1 \rangle| \neq 0 \quad \text{and} \quad |\langle U(\delta_x \otimes v_2), \delta_{y_2} \otimes w_2 \rangle| \neq 0,$$

then $d(y_1, y_2) \leq S$.

2. For all $R > 0$ there exists $S \geq 0$ such that if $x_1, x_2 \in X$ are such that $d(x_1, x_2) \leq R$ and $v_1, v_2 \in \mathcal{H}_1$, $y_1, y_2 \in Y$, and $w_1, w_2 \in \mathcal{H}_1$ are such that

$$|\langle U(\delta_{x_1} \otimes v_1), \delta_{y_1} \otimes w_1 \rangle| \neq 0 \quad \text{and} \quad |\langle U(\delta_{x_2} \otimes v_2), \delta_{y_2} \otimes w_2 \rangle| \neq 0,$$

then $d(y_1, y_2) \leq S$.

The same properties hold with the roles of X and Y reversed, and with U replaced by U^* . □

Proof of Theorem 5.1. Let v_0 denote some fixed choice of unit vector in $\mathcal{H}$. Define a function $f : X \rightarrow Y$ by choosing $f(x)$ to be any element of Y such that there exists $w_x \in \mathcal{H}_1$ such that

$$|\langle U(\delta_x \otimes v_0), \delta_{f(x)} \otimes w_x \rangle| \neq 0.$$

Similarly, define a function $g : Y \rightarrow X$ by choosing $g(y)$ to be any element of X such that there exists $v_y \in \mathcal{H}_1$ such that

$$|\langle U^*(\delta_y \otimes v_0), \delta_{g(y)} \otimes v_y \rangle| \neq 0.$$

The proof concludes exactly as that of Theorem 4.1 above, using Lemma 5.5 in place of Lemma 4.3. □

6 The stable uniform case

In this section we prove our results in the stable uniform case. This is slightly more subtle than the other cases we consider, but essentially the same argument works with a few minor changes. Theorems 4.1, 5.1 and 5.2 above become the following theorem in this context.

Theorem 6.1. *1. Let X and Y be spaces with the metric sparsification property. If there is a $*$ -isomorphism $\phi : C_s^*(X) \rightarrow C_s^*(Y)$, then X and Y are coarsely equivalent.*

2. Let X and Y be spaces with property A. If there is a $*$ -isomorphism $\phi : \mathbb{C}_s[X] \rightarrow \mathbb{C}_s[Y]$, then X and Y are coarsely equivalent.

3. Let X and Y be spaces. If there exists a $*$ -isomorphism $\phi : \mathbb{C}_s[X] \rightarrow \mathbb{C}_s[Y]$ that is spatially implemented by a unitary isomorphism $U : l^2(X, \mathcal{H}) \rightarrow l^2(Y, \mathcal{H})$ (equivalently, that extends to a $*$ -isomorphism $\tilde{\phi} : C_s^*(X) \rightarrow C_s^*(Y)$), then X and Y are coarsely equivalent.

Corollary 6.2. *Let X and Y be spaces with the metric sparsification property. Then $C_u^*(X)$ is Morita equivalent to $C_u^*(Y)$ if and only if X is coarsely equivalent to Y .*

Proof. If X is coarsely equivalent to Y , then $C_u^*(X)$ is Morita equivalent to $C_u^*(Y)$ by [2, Theorem 4]. On the other hand, if $C_u^*(X)$ and $C_u^*(Y)$ are Morita equivalent, then $C_s^*(X) \cong C_s^*(Y)$ by [3, Theorem 1.2], so the result follows from part 1 of Theorem 6.1 above. $\square$

We will concentrate on part 1 of this theorem: parts 2 and 3 follow from the same ingredients, having made essentially the same changes we made when passing from Section 4 to Section 5 above. As in Section 4, fix X and Y with the metric sparsification property and a $*$ -isomorphism ϕ as in the statement of Theorem 6.1, part 1, and let $U : l^2(X, \mathcal{H}) \rightarrow l^2(Y, \mathcal{H})$ be the unitary isomorphism spatially implementing ϕ coming from Lemma 3.1.

The main extra ingredient we need over previous sections is the following lemma.

Lemma 6.3. *Fix a unit vector $v_0 \in \mathcal{H}$ and $c > 0$. Say for each $x \in X$ there exists $f(x) \in Y$ and $w_x \in \mathcal{H}_1$ such that*

$$|\langle U(\delta_x \otimes v_0), \delta_{f(x)} \otimes w_x \rangle| \geq c.$$

Then there exists a finite rank projection $p \in \mathcal{B}(\mathcal{H})$ such that

$$|\langle U(\delta_x \otimes v_0), \delta_{f(x)} \otimes pw_x \rangle| \geq c/2$$

for all $x \in X$. The same statement holds with the roles of X and Y reversed, and with U replaced by U^* .

Proof. We claim first that there exists a finite rank projection $p \in \mathcal{B}(\mathcal{H})$ such that

$$\|Ue_{(x,v_0),(x,v_0)}U^*(1 - 1 \otimes p)\| \leq c/2 \quad (8)$$

for all $x \in X$. Indeed, say otherwise, in which case there exists an increasing sequence of finite rank projections (p_n) in $\mathcal{B}(\mathcal{H})$ that converge strongly to the identity, and a sequence (x_n) of elements of X (which we may assume distinct) such that

$$\|Ue_{(x_n,v_0),(x_n,v_0)}U^*(1 - 1 \otimes p_n)\| \geq c/2$$

for all n . Moreover, there then exist finite rank projections $q_n \in \mathcal{B}(l^2(X, \mathcal{H}))$ such that

$$\|q_n Ue_{(x_n,v_0),(x_n,v_0)}U^*(1 - 1 \otimes p_n)q_n\| \geq c/4$$

for all n . Using finite dimensionality of the projections q_n , an argument based on the Riemann rearrangement theorem as in the proof of Lemma 3.2 shows that there exist subsets $\mathcal{M}, \mathcal{N} \subseteq \mathbb{N}$ such that $\mathcal{N}$ is infinite, and such that

$$\left\| q_n U \left(\sum_{m \in \mathcal{M}} e_{(x_m,v_0),(x_m,v_0)} \right) U^*(1 - 1 \otimes p_n) q_n \right\| \geq c/16$$

for all $n \in \mathcal{N}$. This, however, implies that

$$\left\| U \left(\sum_{m \in \mathcal{M}} e_{(x_m,v_0),(x_m,v_0)} \right) U^*(1 - 1 \otimes p_n) \right\| \geq c/16$$

for all n , which contradicts that

$$U \left(\sum_{m \in \mathcal{M}} e_{(x_m,v_0),(x_m,v_0)} \right) U^* = \phi \left(\sum_{m \in \mathcal{M}} e_{(x_m,v_0),(x_m,v_0)} \right)$$

is an element of $C_s^*(Y)$. This establishes the claim in line (8).

Note then that for any $x \in X$,

$$\begin{aligned} |\langle U(\delta_x \otimes v_0), \delta_{f(x)} \otimes pw_x \rangle| &= \|Ue_{(x,v_0),(x,v_0)}U^*(\delta_{f(x)} \otimes pw_x)\| \\ &\geq \|Ue_{(x,v_0),(x,v_0)}U^*(\delta_{f(x)} \otimes w_x)\| - \|Ue_{(x,v_0),(x,v_0)}U^*(1 - 1 \otimes p)\| \\ &\geq c - c/2 = c/2. \end{aligned}$$

The last statement is clear from symmetry. $\square$

The next lemma follows from the same proof as Lemma 4.3.

Lemma 6.4. *Fix a finite dimensional subspace $\mathcal{V}$ of $\mathcal{H}$.*

1. *Let $\delta > 0$. Then there exists $S \geq 0$ such that if $x \in X$, $v_1, v_2 \in \mathcal{V}_1$, $y_1, y_2 \in Y$ and $w_1, w_2 \in \mathcal{H}_1$ are such that*

$$|\langle U(\delta_x \otimes v_1), \delta_{y_1} \otimes w_1 \rangle| \geq \delta \quad \text{and} \quad |\langle U(\delta_x \otimes v_2), \delta_{y_2} \otimes w_2 \rangle| \geq \delta,$$

then $d(y_1, y_2) \leq S$.

2. Let $\delta > 0$. Then for all $R > 0$ there exists $S \geq 0$ such that if $x_1, x_2 \in X$ are such that $d(x_1, x_2) \leq R$ and $v_1, v_2 \in \mathcal{V}_1$, $y_1, y_2 \in Y$, and $w_1, w_2 \in \mathcal{H}_1$ are such that

$$|\langle U(\delta_{x_1} \otimes v_1), \delta_{y_1} \otimes w_1 \rangle| \geq \delta \quad \text{and} \quad |\langle U(\delta_{x_2} \otimes v_2), \delta_{y_2} \otimes w_2 \rangle| \geq \delta,$$

then $d(y_1, y_2) \leq S$.

The same properties hold with the roles of X and Y reversed, and with U replaced by U^* . $\square$

Proof of Theorem 6.1, part 1. Just as in the proof of Theorem 4.1, let v_0 denote some fixed choice of unit vector in $\mathcal{H}$, and let $c > 0$ have the property in Lemma 4.4 (which still holds in this context, with the same proof) for both X and Y . Define a function $f : X \rightarrow Y$ by choosing $f(x)$ to be any element of Y such that there exists $w_x \in \mathcal{H}_1$ such that

$$|\langle U(\delta_x \otimes v_0), \delta_{f(x)} \otimes w_x \rangle| \geq c$$

(such exists by Lemma 4.4). Similarly define a function $g : Y \rightarrow X$ by choosing $g(y)$ to be any element of X such that there exists $v_y \in \mathcal{H}_1$ such that

$$|\langle U^*(\delta_y \otimes v_0), \delta_{g(y)} \otimes v_y \rangle| \geq c.$$

Using Lemma 6.3 (twice, and taking the supremum of the two projections given), there exists a finite rank projection $p \in \mathcal{B}(\mathcal{H})$ such that

$$|\langle U(\delta_x \otimes v_0), \delta_{f(x)} \otimes pw_x \rangle| \geq c/2$$

for all $x \in X$ and

$$|\langle U^*(\delta_y \otimes v_0), \delta_{g(y)} \otimes pv_y \rangle| \geq c/2$$

for all $y \in Y$.

The proof is then completed just as in the proof of Theorem 4.1, using Lemma 6.4 in place of Lemma 4.3. $\square$

A Categorical interpretation

The aim of this appendix is to provide a slightly more conceptual framework for part 3 of Theorem 1.8. For simplicity, we work throughout with the algebraic Roe algebras $\mathbb{C}[X]$, rather than one of the variants.

We show that coarse equivalences, and spatially implemented isomorphisms of algebraic Roe algebras are ‘essentially the same thing’. More precisely, we define two categories of metric spaces: the morphisms in the first are given by closeness classes of coarse maps, and those in the second by ‘closeness’ classes of spatially implemented isomorphisms of Roe algebras; we then show that these two categories are isomorphic.

It is an unfortunate deficiency of our methods that all morphisms in our categories are isomorphisms: it is not currently clear to us if something more

general is possible. Note also that the main novelty in this section is part 1 of Lemma A.4 (which follows immediately from our techniques in the rest of the piece); the remaining material comes from ideas used by Higson–Roe–Yu in their K -theoretic analyses of Roe algebras (see for example [21, Section 4], which is probably the original reference along these lines.).

Definition A.1. Define a category $\mathcal{C}_0$ by setting the objects to be uniformly discrete, bounded geometry metric spaces, and the morphisms to be coarse equivalences.

Define a category $\mathcal{C}$ to be the quotient category of $\mathcal{C}_0$ under the equivalence relation on morphisms defined by closeness.

Definition A.2. Let X and Y be uniformly discrete, bounded geometry metric spaces, and $U, V : l^2(X, \mathcal{H}) \rightarrow l^2(Y, \mathcal{H})$ be unitary isomorphisms. U and V are said to be *close* if U^*V is a finite propagation operator on $l^2(X, \mathcal{H})$.

Define a category $\mathcal{R}_0$ by setting the objects to be uniformly discrete, bounded geometry metric spaces, and the morphisms to be unitary isomorphisms $U : l^2(X, \mathcal{H}) \rightarrow l^2(Y, \mathcal{H})$ such that $T \mapsto UTU^*$ defines an isomorphism $\mathbb{C}[X] \rightarrow \mathbb{C}[Y]$.

Define a category $\mathcal{R}$ to be the quotient category of $\mathcal{R}_0$ under the equivalence relation on morphisms defined by closeness.

The following definition relates morphisms in $\mathcal{C}_0$ to morphisms in $\mathcal{R}_0$.

Definition A.3. Let $f : X \rightarrow Y$ be a coarse equivalence, and $U : l^2(X, \mathcal{H}) \rightarrow l^2(Y, \mathcal{H})$ a unitary isomorphism. U is said to *cover* f if there exists $C \geq 0$ such that for all $x \in X$ and $y \in Y$, $U_{yx} \neq 0$ implies that $d(f(x), y) \leq C$.

Lemma A.4. 1. For any morphism U in $\mathcal{R}_0$, there exists a morphism f in $\mathcal{C}_0$ covered by U .

2. For any morphism f in $\mathcal{C}_0$, there exists a morphism U in $\mathcal{R}_0$ that covers f .

Moreover, any two morphisms in $\mathcal{R}_0$ covering the same morphism in $\mathcal{C}_0$ are close, and any two morphisms in $\mathcal{C}_0$ covered by a single morphism in $\mathcal{R}_0$ are close.

Proof. For the first part, let f be built from U as in the proof of Theorem 5.1; clearly U covers f .

The second part is folklore, but does not seem to appear in the literature: this is perhaps due to K -theoretic arguments requiring only covering *isometries*, not covering *unitaries*. We sketch a proof. Let $f : X \rightarrow Y$ be a morphism in $\mathcal{C}_0$. Consider any partition $Y = \sqcup Y_n$ with the properties:

- there exists $C > 0$ such that $\text{diam}(Y_n) \leq C$ for all n ;
- for all n , $Y_n \cap \text{Image}(f) \neq \emptyset$.

Using the fact that f is a coarse equivalence, it is not difficult to see that such a partition exists. Define a partition $X = \sqcup X_n$ by setting $X_n = f^{-1}(Y_n)$, and fix arbitrary unitary isomorphisms $U_n : l^2(X_n, \mathcal{H}) \rightarrow l^2(Y_n, \mathcal{H})$ (such exist as the domain and codomain are separable infinite dimensional Hilbert spaces). Define U to be the unitary

$$U := \oplus U_n : l^2(X, \mathcal{H}) \rightarrow l^2(Y, \mathcal{H});$$

it is not hard to see that U is a morphism in $\mathcal{R}_0$ that covers f .

The final comments follow from simple computations. $\square$

Given the preceding lemma, the proof of the following theorem is a series of routine checks, and is thus omitted.

Theorem A.5. *Provisionally define functors*

$$\mathcal{F} : \mathcal{R} \rightarrow \mathcal{C} \quad \text{and} \quad \mathcal{U} : \mathcal{C} \rightarrow \mathcal{R}$$

as follows: each is the identity on objects; $\mathcal{F}[U]$ is the class in $\mathcal{C}$ of any coarse equivalence covered by U ; $\mathcal{U}[f]$ is the class in $\mathcal{R}$ of any unitary isomorphism covering f .

Then $\mathcal{F}$ and $\mathcal{U}$ are well-defined and mutually inverse isomorphisms of categories. $\square$

Acknowledgements

This piece grew out of questions asked (independently) of us by our doctoral advisors: Guoliang Yu for Špakula and John Roe for Willett. We are very grateful to Yu and Roe for suggesting this line of research, and for their ongoing encouragement and support.

An earlier version of the piece used finite asymptotic dimension where we currently use the much more flexible metric sparsification property. We would like to thank Romain Tessera for suggesting the use of metric sparsification.

The first author was supported by the *Deutsche Forschungsgemeinschaft* (SFB 878).

References

- [1] G. Arzhantseva and T. Delzant. Examples of random groups. Available on the authors' websites, 2008.
- [2] J. Brodzki, G. Niblo, and N. Wright. Property A, partial translation structures and uniform embeddings in groups. *J. London Math. Soc.*, 76(2):479–497, 2007.
- [3] L. G. Brown, P. Green, and M. A. Rieffel. Stable isomorphism and strong morita equivalence of C^* -algebras. *Pacific J. Math.*, 71(2):349–363, 1977.

- [4] N. Brown and N. Ozawa. *C*-Algebras and Finite-Dimensional Approximations*, volume 88 of *Graduate Studies in Mathematics*. American Mathematical Society, 2008.
- [5] X. Chen, R. Tessera, X. Wang, and G. Yu. Metric sparsification and operator norm localization. *Adv. Math.*, 218(5):1496–1511, 2008.
- [6] X. Chen and Q. Wang. Ideal structure of uniform Roe algebras of coarse spaces. *J. Funct. Anal.*, 216(1):191–211, 2004.
- [7] X. Chen and Q. Wang. Ghost ideals in uniform Roe algebras of coarse spaces. *Arch. Math. (Basel)*, 84(6):519–526, 2005.
- [8] A. Connes. *Noncommutative Geometry*. Academic Press, 1994.
- [9] J. Dixmier. *C*-Algebras*. North Holland Publishing Company, 1977.
- [10] G. Gong, Q. Wang, and G. Yu. Geometrization of the strong Novikov conjecture for residually finite groups. *J. Reine Angew. Math.*, 621:159–189, 2008.
- [11] M. Gromov. Asymptotic invariants of infinite groups. In G. Niblo and M. Roller, editors, *Geometric Group Theory*, volume 2, 1993.
- [12] M. Gromov. Random walks in random groups. *Geom. Funct. Anal.*, 13(1):73–146, 2003.
- [13] E. Guentner, N. Higson, and S. Weinberger. The Novikov conjecture for linear groups. *Publ. Math. Inst. Hautes Études Sci.*, 101:243–268, 2005.
- [14] E. Guentner and J. Kaminker. Exactness and the Novikov conjecture. *Topology*, 41(2):411–418, 2002.
- [15] E. Guentner, R. Tessera, and G. Yu. A notion of geometric complexity and its application to topological rigidity. Available on the arxiv, 2009.
- [16] E. Guentner, R. Tessera, and G. Yu. Operator norm localization for linear groups and its applications to K -theory. *Adv. Math.*, 226(4):3495–3510, 2010.
- [17] N. Higson. Counterexamples to the coarse Baum-Connes conjecture. Available on the author’s website, 1999.
- [18] N. Higson, V. Lafforgue, and G. Skandalis. Counterexamples to the Baum-Connes conjecture. *Geom. Funct. Anal.*, 12:330–354, 2002.
- [19] N. Higson and J. Roe. On the coarse Baum-Connes conjecture. *London Mathematical Society Lecture Notes*, 227:227–254, 1995.
- [20] N. Higson and J. Roe. Amenable group actions and the Novikov conjecture. *J. Reine Angew. Math.*, 519:143–153, 2000.

- [21] N. Higson, J. Roe, and G. Yu. A coarse Mayer-Vietoris principle. *Math. Proc. Cambridge Philos. Soc.*, 114:85–97, 1993.
- [22] A. Ioana. W^* -superrigidity for Bernoulli actions of property (T) groups. Available on the author’s website, 2010.
- [23] N. Ozawa. Amenable actions and exactness for discrete groups. *C. R. Acad. Sci. Paris Sér. I Math.*, 330:691–695, 2000.
- [24] S. Popa. Deformation and rigidity for group actions and von Neumann algebras. In *Proceedings of the International Congress of Mathematicians*, volume I, pages 445–477. Eur. Math. Soc., 2007.
- [25] J. Roe. An index theorem on open manifolds, I. *J. Differential Geometry*, 27:87–113, 1988.
- [26] J. Roe. Coarse cohomology and index theory on complete Riemannian manifolds. *Mem. Amer. Math. Soc.*, 104(497), 1993.
- [27] J. Roe. *Index Theory, Coarse Geometry and Topology of Manifolds*, volume 90 of *CBMS Conference Proceedings*. American Mathematical Society, 1996.
- [28] J. Roe. *Lectures on Coarse Geometry*, volume 31 of *University Lecture Series*. American Mathematical Society, 2003.
- [29] J. Roe. Hyperbolic groups have finite asymptotic dimension. *Proceedings of the American Mathematical Society*, 133(9):2489–2490, 2005.
- [30] J. Špakula. Non- K -exact uniform Roe C^* -algebras. To appear in *Journal of K -theory*; available on the author’s website, 2009.
- [31] J. Špakula. Uniform K -homology theory. *J. Funct. Anal.*, 257(1):88–121, 2009.
- [32] J. Špakula and R. Willett. Maximal and reduced Roe algebras of coarsely embeddable spaces. Accepted to *J. Reine Angew. Math.*; available on the authors’ websites, 2010.
- [33] Q. Wang. Remarks on ghost projections and ideals in the Roe algebras of expander sequences. *Arch. Math. (Basel)*, 89(5):459–465, 2007.
- [34] R. Willett. Some notes on property A. In G. Arzhantseva and A. Valette, editors, *Limits of Graphs in Group Theory and Computer Science*, pages 191–281. EPFL press, 2009.
- [35] W. Winter and J. Zacharias. The nuclear dimension of C^* -algebras. *Adv. Math.*, 224(2):461–498, 2010.
- [36] G. Yu. Coarse Baum-Connes conjecture. *K-theory*, 9:199–221, 1995.

- [37] G. Yu. The Novikov conjecture for groups with finite asymptotic dimension.
Ann. of Math., 147(2):325–355, 1998.
- [38] G. Yu. The coarse Baum-Connes conjecture for spaces which admit a uniform embedding into Hilbert space. *Invent. Math.*, 139(1):201–240, 2000.

MATHEMATISCHES INSTITUT, UNIVERSITÄT MÜNSTER, EINSTEINSTR. 62, 48149
MÜNSTER, GERMANY
E-mail: `jan.spakula@uni-muenster.de`

2565 MCCARTHY MALL, UNIVERSITY OF HAWAI‘I AT MĀNOA, HONOLULU,
HI 96822, USA and
1326 STEVENSON CENTER, VANDERBILT UNIVERSITY, NASHVILLE, TN 37240,
USA
E-mail: `rufus.willet@vanderbilt.edu`